



Epidemiological, Characteristics and Complication of Clavicular Fractures among Patients in Baghdad

Saman Mohammad AbdulRazaq*

Corresponding Author: Saman Mohammad AbdulRazaq, MBCHB, FIBMS, FABMS. Baghdad /Iraq.

Copy Right: © 2021 Dr Suresh Kumar N Parmar. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received Date: June 17, 2021

Published date: July 01, 2021

Abstract

In recent years, collarbone fractures have been a common occurrence among adults and the elderly. This study aimed to assess the characteristics of clavicle fracture among patients and, also to identify the complication of clavicle fracture; whether it had an impact on treatment ways during the period of study. A cross-sectional study was conducted in eight general hospitals in Baghdad during the period from August 2020 up to May 2021 to determine the epidemiology and characteristics of people with collarbone fractures. The sample size was 130 samples. The information was collected directly from the patient or the patient's relative. STATA 20 software package was used to calculate the figures and perform all the tests at the P-value <0.05 was considered as statistically significant. The mean age and SD were 38.4±14.3; the highest frequency 30% of clavicle fracture occurred among them in the age groups more than 45 years old and 27.69% in the age 26 to 35 years old. The majority of them were male 75.38% and 24.62% were female. Most of the 71.54% were exposed to clavicular fracture as an accident by car, cycling and motorcycle. 59.23% of them suffering from a middle fracture.

There is highly significant association has been found between the age groups and AO/OTA classification at the p. value =0.0002 and, a significant relationship has been found between the sites of fracture with AO/OTA classification at the p. value less than 0.05. Weak /slow recoveries were the main complication of clavicle fracture among studied samples. We need to encourage clinicians to attend scientific conferences and educational seminars.

Keyword: - Fracture, Clavicle, Accident, Complication, Assess, Baghdad.

Introduction

Clavicle fractures occur commonly, often as a result of indirect or direct trauma to the shoulder region [1, 2]. The incidence of clavicle fractures in the adolescent and adult population is suggested to be between 29 and 64 per 100.000 persons [3]. As usual, in many traumas, its prevalence is highest among the young population even if also shows a bimodal age distribution with a rate in females that overtake males after the sixth decade of life as a result of osteoporosis and differences in life expectancy [4,5]. The mean age has been reported to be 29.3 years, and the incidence appears to decrease significantly after the second decade of life [6]. Males are affected approximately twice as often as females (67.9% vs 32.1%) [6]; these injuries may also have a seasonal correlation, with one epidemiologic analysis noting an increase during the summer [6, 7]. In adults, more than two-thirds of these injuries occur at the diaphysis of the clavicle, and these injuries are more likely to be displaced as compared with medial and lateral third fractures [8,9]. Lateral-third fractures are less common, accounting for approximately 25% of all clavicle fractures, and are less likely to be displaced than those occurring in the midshaft[10,11]. Medial-third fractures comprise the remaining 2% to 3% of these injuries [12]. In Iraq, there is little research on this field because of the lack of the necessary capabilities to conduct such research, in addition to, the deterioration of the health situation and the spread of ignorance about the complications of the fracture and the extent of its future impact on their health. Also, surgeons turned to the old methods of treatment due to the lack of ways to develop capabilities and hold seminars that help researchers to share and develop their ideas from the point we need to encourage the clinicians who are working in emergency departments and general clinics and should be familiar with the common presentations and complications of this injury, as well as basic management. Despite the high frequency, the choice of proper treatment is still a challenge for orthopedic surgeons.

This study aimed to assess the characteristics of clavicle fracture among patients and, also to identify the complication of clavicle fracture; whether it had an impact on treatment ways during the period of study.

Methodology

A cross-sectional study was conducted in eight general hospitals in Baghdad during the period from August 2020 up to May 2021 in order to determine the epidemiology and characteristics of people with collarbone fractures. The sample size was 130 samples. Ethical clearance was obtained from the Iraqi Ministry of Health before collecting the data. The information was collected directly from the patient or the patient's relative in the event that he /she was unable to answer the questionnaire, and then the data was checked with the hospital records, and if there was any difference, the case was excluded from the study. The inclusion criteria were patients aged 16 or greater with fresh clavicular fractures. The exclusion criteria were patients under 16 years, those with only ligamentous injuries or dislocation; old fractures and clavicular fractures due to neoplasm metastasis or other medical conditions. Details of all the patients regarding their age, gender, involved side and injury date were recorded. AO/OTA classification was firstly reported and evolved for better clinical research and surgical decision-making for adult fracture (age >16 years) [13]. The details on this classification for clavicular fracture were as follows: "15" for its anatomical location and three groups were divided as proximal (15A), shaft (15B), and distal (15C) according to the Heim Square method. Continuous variable (age) was expressed as mean and standard deviation and the test was performed using Student t-test or Whitney U-test, based on the normality status of data. Discontinuous variables regarding age (dichotomy), gender, side involved and seasonal variation were expressed as count and percentage and the results were analyzed by Chi-square test. A $P < 0.05$ was considered statistically significant. STATA 20 software package was used to calculate the figures and perform all the tests.

Results

Out of one hundred and thirty cases with clavicular fractures, the mean age and SD were 38.4 ± 14.3 . The highest frequency 30% of clavicle fracture occurred among them in the age groups more than 45 years old and 27.69% in the age 26 to 35 years old. The majority of them were male 75.38% and 24.62% were female. Most of the 71.54% were exposed to clavicular fracture as an accident by car, cycling and motorcycle. 59.23% of them suffering from a middle fracture. The highest frequency 34.62% of cases was diaphysis regarding AO/OTA classification. The highest frequency 36.92% of clavicular fracture cases occurred during the summer season more than other seasons [Table 1]. There is highly significant association has been found between the age groups and AO/OTA classification at the p. value = 0.0002. Also, there is a significant relationship has been found between the sites of fracture with AO/OTA classification at the p. value less than 0.05 [Table 2]. In the same table 2, there is not a significant relationship has been found between gender and causes by AO classification at the p. value of more than 0.05. Distribution of clavicle fracture among studied samples by season [Figure 1]. AO/OTA

Classification of clavicular fractures among studied samples [Figure2]. Weak /slow recovery of 56.2% was the main complication of clavicle fracture [Figure3].

Table 1: Characteristics of studied samples (130)

Variables			
Age	Mean ±Sd	38.4±14.3	
		Number	%
Age groups years	16-25	30	23.08
	26-35	36	27.69
	36-45	25	19.23
	>45	39	30.00
Gender	Male	98	75.38
	Female	32	24.62
Causes	Car, Cycling, Motorcycle	93	71.54
	Soccer	29	22.31
	Heavy & light things	8	6.15
Site of Fracture	Type I (Middle)	77	59.23
	Type II (Acromial)	35	26.92
	Type III (Sternal)	18	13.85
AO/OTA classification	15A (Proximal)	44	33.85
	15B (Diaphysis)	45	34.62
	15C(Distal)	41	31.54
Seasonal	Spring	23	17.69
	Summer	48	36.92

	Autumn	30	23.08
	Winter	29	22.31

Table 2: Distribution of studied samples with clavicle fracture according to age groups by AO/OTA classification

Age groups	AO//OTA classification						Total	P. value	
	15A(44)		15B(45)		15C(41)				
16-25	13	29.55	13	28.89	4	9.76	30	23.08	0.0002
26-35	18	40.91	13	28.89	5	12.20	36	27.69	
36-45	9	20.45	9	20.00	7	17.07	25	19.23	
>45	4	9.09	10	22.22	25	60.98	39	30.00	
Gender									
Male	32	72.73	32	71.11	34	82.93	98	75.38	0.3931
Female	12	27.27	13	28.89	7	17.07	32	24.62	
Causes									
Car, cycling, motorcycle	30	68.18	29	64.44	34	82.93	93	71.54	0.2234
Soccer	12	27.27	11	24.44	6	14.63	29	22.31	
Heavy & light things	2	4.55	5	11.11	1	2.44	8	6.15	
Site of fracture									
Middle	32	72.73	28	62.22	17	41.46	77	59.23	0.0049
Acromial	4	9.09	14	31.11	17	41.46	35	26.92	
Sternal	8	18.18	3	6.67	7	17.07	18	13.85	




Figure 1: Distribution of clavicle fracture among studied sample by season




Figure 2: AO/OTA Classification of clavicular fractures




Figure 3: Complication of clavicle fracture among patients

Discussion

This study aimed to assess the characteristics of clavicle fracture among patients and, also to identify the complication of clavicle fracture; whether it had an impact on treatment ways during the period of study. The clavicle fracture doesn't become fully solid until about age 20, which makes teens more likely to fracture a collarbone [14]. The risk decreases after age 20, but rises again in older adults, as bone strength declines with age [15]. In our finding shows that the mean age was 38.4 year with Sd 14.3 year compared with other studied done it in Sweden by Kihlström, et al, the authors found the mean age was higher in females (mean 59 years, SD 23 years) than in males (mean 43 years, SD 21 years)[16]. While a study was done in Italy by Paladini, et al, they found the mean age was 29.3 years [17], this refers to the difference in the lifestyle between the two countries, in addition to, the physical condition of each person, for example, in developed countries, the elderly suffer from chronic diseases and aging, on the contrary in developing countries, where young people and the elderly suffer from various diseases according to the conditions. Also, we found 30% of cases with clavicular fracture occurred in the age group more than 45 years old and compared this result with others in Sweden by Nowak [18], they reported the majority were in the age 25 to 35 years old. A study in the US [19], found the most of them were in the age above 40 years old. This refers to the difference in the type of activity which is practicing by them. Clavicular fractures are most common in young adults and elderly women [20]. In this study, we found the majority 75.38% of clavicular fracture patients occurred among men more than females.

A study did it in Ireland [21] and in the US [22], the authors found the men are more likely to get a clavicle fracture than females. This explains that most of the hard work is practiced by men, in addition to, playing sports such as football, swimming and weight lifting. Clavicle fractures are a common injury, especially in children and young adults. The collarbone connects the top of the sternum to the scapula, and common causes of broken collarbones include falls, sports injuries, and traffic accident trauma [23, 24]. Car, cycling and motorcycle are the main causes of clavicular fracture among cases, and it has been showing in 71.54% and compared with other studied done it in the US [22, 23], Africa [24], China [25], the investigators found the car and motorcycle is the major type of traffic accident. This similarity is due to the use of the same means of transportation, which is a major cause of injury, in addition to, the ignorance of most of them about safety and security conditions when they are using means of transportation. In our study, we found 59.23% of them were suffering from middle fracture and compared with other studied in Turkey [26] and in the US [27]. They found the majority are suffering from a sternal and acromial fracture. This depends on the type of external pressure and how strongly it affects the body. In this study, we found the most cases 36.92% occurred during the summer season and in a study in the US [28], the authors found the most of the fractures occurred during the autumn & winter seasons. This refers to the difference in lifestyle between countries, as well as the ways to use transportation, especially during the winter season. Clavicle fractures in adolescents may take six to eight weeks. However, fractures in adults or teenagers who have stopped growing take 10 to 12 weeks to heal and may take longer [29]. Also, we found 56.2% of cases were suffering from weak or slow recovery from clavicular fractures and compared with another study in Italy [17] and in India [30]. This depends on the ways of treatment which are used by the physicians.

Conclusion

we concluded the clavicle fracture was common among men. The highest frequency of clavicle fracture occurred among age groups more than 45 years old. Car, cycling and motors are the main cause of fracture. The summer season was a higher rate more than other seasons. The middle site of fracture was the highest frequency among others. There is highly significant association has been found between the age groups and AO/OTA classification at the p. value =0.0002 and, a significant relationship has been found between the sites of fracture with AO/OTA classification at the p. value less than 0.05. In addition, there are not significant relationship has been found between gender and causes by AO classification at the p. value of more than 0.05. Weak /slow recoveries were the main complication of clavicle fracture.

Recommendation: we recommended encouraging all the clinicians to attend scientific conferences and

educational seminars that promote use the of modern treatment ways and to see the latest findings of experts in the orthopedics field.

Conflict of interest: Nill

References

1. Ropars M, Thomazeau H, Hutten D (February 2017). "Clavicle fractures". *Orthopaedics & Traumatology, Surgery & Research*. 103 (1S): S53–S59
2. Postacchini F, Gumina S, De Santis P, Albo F. "Epidemiology of clavicle fractures". *J Shoulder Elbow Surg*. 2002;11(5):452–6.
3. Huttunen TT, Launonen AP, Berg HE, Lepola V, Fellander-Tsai L, Mattila VM. "Trends in the Incidence of Clavicle Fractures and Surgical Repair in Sweden: 2001–2012". *J Bone Joint Surg Am*. 2016;98(21):1837–42.
4. Toogood P, Horst P, Samagh S, Feeley BT. "Clavicle fractures: a review of the literature and update on treatment". *Phys Sportsmed*. 2011 Oct;39(3):142–50
5. Khan LA, Bradnock TJ, Scott C, Robinson CM. "Fractures of the clavicle". *J Bone Joint Surg Am*. 2009 Feb;91(2):447–60.
6. Amer KM, Congiusta DV, Suri P, Choudhry A, Otero K, Adams M. "Clavicle fractures: Associated trauma and morbidity". *J Clin Orthop Trauma*. 2020 Aug 26;13:53-56.
7. Alao D, Guly HR. "Missed clavicular fracture; inadequate radiograph or occult fracture?" *Emerg Med J*. 2005;22 (3): 232-3.
8. Labronici PJ, Santos Filho FCD, Reis TB, Pires RES, Junior AFM, Kojima KE. "Are diaphyseal clavicular fractures still treated traditionally in a non-surgical way?" *Rev Bras Ortop*. 2017 Jul 8;52(4):410-416.
9. Asadollahi S, Bucknill A. "Acute medial clavicle fracture in adults: a systematic review of demographics, clinical features and treatment outcomes in 220 patients". *J Orthop Traumatol*. 2019 Jun 28;20(1):24.
10. Hoogervorst P, van Schie P, van den Bekerom MP. "Midshaft clavicle fractures: Current concepts". *EFORT Open Rev*. 2018 Jun 20;3(6):374-380.
11. Jeray KJ. "Acute midshaft clavicular fracture". *J Am Acad Orthop Surg*. 2007 Apr;15(4):239–48.

12. Throckmorton T, Kuhn JE. "Fractures of the medial end of the clavicle". *J Shoulder Elbow Surg.* 2007;16(1):49-54.
13. Meinberg EG, Agel J, Roberts CS, Karam MD, Kellam JF. "Fracture and Dislocation Classification Compendium-2018". (2018) *Journal of orthopaedic trauma.* 32 Suppl 1: S1-S170.
14. Biglari F, Mavaein A, Shabani S, Mahdavi Mohtasham H, Kazemi S M. "Complication of Clavicular Fractures after Open Reduction". *IJOS.* 2017; 15 (2) :39-45.
15. Millett PJ, Hurst JM, Horan MP, Hawkins RJ. "Complications of clavicle fractures treated with intramedullary fixation". *J Shoulder Elbow Surg.* 2011 Jan;20(1):86–91. Epub 2010 Nov 3.
16. Kihlström, C., Möller, M., Lönn, K. et al. "Clavicle fractures: epidemiology, classification and treatment of 2 422 fractures in the Swedish Fracture Register; an observational study". *BMC Musculoskelet Disord* 18, 82 (2017).
17. Paladini P, Pellegrini A, Merolla G, Campi F, Porcellini G. "Treatment of clavicle fractures". *Transl Med UniSa.* 2012 Jan 18;2:47-58.
18. Nowak J, Mallmin H, Larsson S. The aetiology and epidemiology of clavicular fractures. "A prospective study during a 2-year period in Uppsala, Sweden". *Injury.* 2000;31(5):353–8.
19. Hsiao MS, Cameron KL, Huh J, Hsu JR, Benigni M, Whitener JC, Owens BD. "Clavicle fractures in the United States military: incidence and characteristics". *Mil Med.* 2012 Aug;177(8):970-4.
20. Dyan V. Flores, Paola Kuenzer Goes, Catalina Mejía Gómez, Darwin Fernández Umpire, Mini N. Pathria. "Imaging of the Acromio clavicular Joint: Anatomy, Function, Pathologic Features, and Treatment". (2020) *RadioGraphics.* 40 (5): 1355-1382.
21. O'Neill BJ, Hirpara KM, O'Briain D, McGarr C, Kaar TK. "Clavicle fractures: a comparison of five classification systems and their relationship to treatment outcomes". *Int Orthop.* 2011 Jun;35(6):909–14. Epub 2010 Nov 19.
22. Luo TD, Ashraf A, Larson AN, Stans AA, Shaughnessy WJ, McIntosh AL. "Complications in the treatment of adolescent clavicle fractures". *Orthopedics.* 2015;38(4):e287-e91.
23. Yang, S, Werner, BC, Gwathmey, FW. "Treatment trends in adolescent clavicle fractures". *J Pediatr Orthop.* 2015;35(3):229–233.
24. S Mohammedali, SK Mutiso, P Oroko, H Saidi. Experience with treatment of clavicle fractures at an African Tertiary Referral Hospital. *East African Orthopaedic Journal.* Vol.7: March 2013.

25. Zhu X, Li W, Chen Z. "Analyses of epidemiology in 363 cases of clavicle fractures". *Zhongguo Xiu Fu Chong Jian Wai Ke Za Zhi*. 2004 Jul;18(4):275-6.
26. Kabak S, Halici M, Tuncel M, Avsarogullari L, Karaoglu S. "Treatment of midclavicular nonunion: comparison of dynamic compression plating and low-contact dynamic compression plating techniques" *J Shoulder Elbow Surg* 2004 July-Aug;13(4):396-403.
27. Fanter NJ, Kenny RM, Baker CL 3rd, Baker CL Jr. "Surgical treatment of clavicle fractures in the adolescent athlete". *Sports Health*. 2015 Mar;7(2):137-41
28. Pecci M. & Kreher, J. "Clavicle Fractures" *Am Fam Physician*. 2008 Jan 1;77(1):65-70.
29. Simon RR, Sherman SC, Koenigsknecht SJ. Clavicle fractures. "In *Emergency Orthopaedics - The Extremities*". 5th Ed. McGraw-Hill, Chicago 2007. P.285-7.
30. Lakhotia D, Khatri K, Sharma V, Farooque K, Sharma S. "Bilateral Clavicle Fractures: *Bilateral Clavicle Fractures: A Report of Three Cases*.2016"; 10(6), RR01-RR03.