



Thrombus In Transit.

A Case Report of Right Ventricular Thrombus with Massive Pulmonary Embolism and Right Ventricular Strain.

Dr. Zahid Khan*, Dr Ayub Khan¹, Dr Gideon Mlawa², Dr Mildred Ibekwe³, Dr Shah Nawaz⁴

1,2,3,4, Barking, Redbridge and Havering NHS Health Trust.

Corresponding Author: Dr Zahid Khan, Department of Cardiology, Barking, Redbridge, and Havering NHS Health Trust, United Kingdom.

Copy Right: © 2021 Dr Zahid Khan. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Received Date: October 25, 2021

Published Date: November 09, 2021

Summary

We present a case of 72-year-old patient who presented with chest pain, shortness of breath (sob) for 2 days and 3 episodes of collapses with loss of consciousness (LOC). His past medical history includes Prostate carcinoma, hypercholesterolemia and he had AstraZeneca covid vaccines 6 months ago. He was hemodynamically stable and had urgent computerized tomography pulmonary angiography (CTPA), confirming bilateral pulmonary embolism with evidence of right heart strain. His clinical examination was unremarkable apart from dyspnoea. He was commenced on twice-daily treatment dose Clexane and was not thrombolysed in view of hemodynamic stability. He had urgent bedside echocardiogram confirming right ventricular thrombus with associated right heart strain, with no evidence of patent foramen ovale (PFO). Electrocardiogram (ECG) showed right bundle branch block (RBBB) and left axis deviation (LAD). He was discharged home after three weeks on Rivaroxaban 15 mg BD for 21 days followed by 20 mg OD.

Background

Right heart thrombi (RVT) are uncommon, usually found in association with pulmonary embolism (PE) and are mostly in transit originating from deep venous thrombosis (DVT) in absence of atrial fibrillation (AF). It can occasionally arise from AF and heart failure. The current treatment strategy for RHT and PE is unclear as there have been no randomized controlled trials (RCTs) evaluating various treatments. The incidence of right ventricle thrombus associated with PE is about 4% to 18% and its presence indicates increased mortality risk compared to PE alone [1]. The mortality associated with RVT associated with PE has been reported as 44% and 27% [2,3]. Due to lack of research, there is no clear consensus on the treatment of these patients and various treatment strategies such as anticoagulation, thrombolysis and embolectomy have been used. The diagnostic modality of choice for these patients is computerized tomography pulmonary angiography (CTPA) however this can be difficult particularly in patients with renal failure and hemodynamically instability.

Case Presentation

A 72-year-old patient presented with two days history of SOB and 3 collapses with LOC. His vitals were heart rate 91 beats per minute (bpm), blood pressure (BP) 156/86, capillary refill time (CRT) 3 seconds, respiratory rate 19, afebrile and saturation were 95% on 60% Venturi mask. His COVID 19 lateral flow was negative and the polymerase chain reaction test was also negative. His clinical examination was normal and there was no calf tenderness to suggest deep vein thrombosis (DVT). His blood showed mildly raised white cell count 12.8, neutrophil 9.2, creatinine 122, C-reactive protein 10 and D-Dimer was 23.18 (table 1). Arterial blood gas (ABG) showed type 1 respiratory failure with PO₂ 8.4, PCO₂ 4.4, normal Ph and lactate. ECG showed right bundle branch block (RBBB), left axis deviation (LAD) and sinus rhythm (figure 1). He weighed 89.6 kilograms (kg) and was commenced on a twice-daily treatment dose of low molecular weight heparin (LMWH).

He had urgent CTPA, confirming extensive bilateral PEs with evidence of right heart strain (figure 2 & 3). The patient was discussed with ITU and respiratory consultant on call for possible thrombolysis however a Page 5 of 10 decision was made against it due to his hemodynamic stability. The patient was also discussed with an interventional radiologist for consideration of embolectomy. The patient had a bedside echocardiogram to assess for PFO that confirmed the presence of right ventricular thrombus and dilated right ventricle. He also had a departmental echocardiogram that showed the normal left ventricular size and systolic function with septal flattening suggestive of right heart overload and dilated Right Heart with impaired radial and preserved longitudinal function (Video 1, 2). The pulmonary pressure was elevated with estimated pulmonary artery systolic pressure (PASp) = 66mmHg + right atrial pressure (RAp). Echocardiogram also showed an echogenic mobile mass in the right ventricle without any evidence of PFO (Figure 4). The patient had computerized tomography of the neck and thorax, abdomen

and pelvis that did not show any evidence of malignancy. He had a previous history of prostate cancer which was under remission and the CT scan showed only a mildly enlarged prostate. The autoimmune screen was negative.

He did not have any history of weight loss, was a lifelong non-smoker and received both doses of AstraZeneca vaccine 6 months ago. He was in the hospital for 3 weeks and made a complete recovery.

Investigations

Blood test	Reference range	Day 1	Day 10
Haemoglobin	133 - 173 g/L	141	126
White cell count	3.8 - 11 10 ⁹ /L	12.8	11.7
Neutrophil	2 - 7.5 10 ⁹ /L	9.2	8.2
Platelet	150 - 400 10 ⁹ /L	101	295
Urea	2.5 - 7.8 mmol/L	6.9	8.2
Creatinine	59 - 104 umol/L	122	114
Sodium	133 - 146 mmol/L	141	137
Potassium	3.5 - 5.3 mmol/L	4.2	4.4
C reactive protein	0 - 5 mg/L	10	142
D-Dimer	0 - 0.5 mg/L FEU	23.18	Test not repeated
2019-Novel Coronavirus	SARS-COV-2 RNA not detected	2019-Novel Coronavirus	SARS-COV-2 RNA not detected
Troponin T	0-14	75	Test not repeated

Table 1

Figure/Video Captions




Figure 1. ECG shows RBBB, LAD and sinus rhythm

Citation: Dr Zahid Khan. "Thrombus In Transit. A Case Report of Right Ventricular Thrombus with Massive Pulmonary Embolism and Right Ventricular Strain." MAR Cardiology 3.5
www.medicalandresearch.com (pg. 4)



Figure 2 & 3. CTPA shows extensive PE extending up to the right ventricle.



Video 1. Parasternal long axis showing RV thrombus



Video 2. Apical 4 chambers views showing RV thrombus and dilated right ventricle.

Differential Diagnosis

If Relevant

The possible differential diagnosis include infection, pneumothorax, pleural effusion, pulmonary embolism, pulmonary oedema secondary to myocardial infarction (MI), or myocarditis.

The inflammatory markers were only marginally raised however his clinical examination was unremarkable and he was afebrile. The CTPA and other tests also did not show any evidence of infection.

There was no pleural effusion on CTPA, CTAP and echocardiogram also did not show any evidence of pleural effusion.

This patient had extensive Pulmonary embolism and RV thrombus confirmed by CTPA and echocardiogram.

Patients with pulmonary oedema can present with low oxygen saturation however his chest examination was clear. Additionally, there was no evidence of pulmonary oedema on chest x-ray and CTPA. His

Page 6 of 10 troponin T was 75 but there were no ischaemic changes on his ECG and raised troponin T was due to extensive PEs. He did not have a recent infection and had no ECG features suggestive of myocarditis.

Treatment

If Relevant

This patient was admitted with 2 days history of SOB and 3 collapses with LOC. His blood gases showed type 1 respiratory failure and was requiring 60% oxygen via venturi mask to maintain oxygen saturation above 94%. He was commenced on 90 mg clexane twice daily and was not thrombolysed given his hemodynamic stability. His oxygen requirement reduced over the next few days and was weaned off oxygen. He remained afebrile throughout the admission and was maintaining saturations above 95% without oxygen requirement. He was discharged home on Rivaroxaban 15 mg BD for 21 days followed by 20 mg OD with a 1-month outpatient follow-up. Repeat echocardiogram before discharge showed a reduction in RV size and the size of thrombus.

Outcome and Follow-Up

This patient was followed up in respiratory outpatient clinic two months after discharge and showed good resolution of symptoms. He was also seen by a Haematology consultant in the outpatient clinic

Citation: Dr Zahid Khan. "Thrombus In Transit. A Case Report of Right Ventricular Thrombus with Massive Pulmonary Embolism and Right Ventricular Strain." MAR Cardiology 3.5
www.medicalandresearch.com (pg. 7)

and was happy with his progress. He also had a repeat echo as an outpatient and showed complete resolution of the thrombus.

Discussion

Include a very brief review of similar published cases

RV thrombus and pulmonary embolism can occur together and carries significant mortality risk. There is a lack of research in this regard and there is a lack of guidelines for treating such cases. Most cases of RV thrombus and PEs are due to atrial fibrillation or patent foramen ovale, however, there was no evidence of either in this case. Malignancy is another possible cause for extensive pulmonary embolism however there was no evidence of malignancy in his case.

The European Working Group on ECG has identified three patterns of the right heart thrombi based on 2D echocardiogram findings.[5] Type A thrombi are highly mobile and are likely dislodged from peripheral venous clots. Patients with type A clots have a high mortality risk of up to 42% due to extreme mobility of the clots. Type B thrombi are similar to left heart thrombi, are almost immobile indicating their in-situ formation and their mortality risk is about 4%. The third type Type C thrombi are rare, share characteristics of both type A and C and are highly mobile. They resemble myxoma in appearance and have a mortality rate intermediate between the above two types. The echocardiogram in our case showed a mobile echogenic mass in the right heart traversing through the tricuspid valve measuring 2cm² indicating type C thrombus and there was no evidence of peripheral venous thrombosis.

There is no clear consensus regarding the treatment of right heart thrombus in transit with extensive pulmonary embolism and treatment options include anticoagulation, thrombolysis, and embolectomy. The success rate of treatment depends on patient clinical status and Rose et al. reported the mortality rate as 100%, 28.6%, 23.8%, 11.3% with no therapy, anticoagulation, surgical embolectomy, and thrombolysis. However, no significant difference between these treatment options was reported in other studies based on in-hospital mortality [6,7,8].

Another case study of a 63-year-old patient with right heart thrombus and massive pulmonary embolism was successfully treated with apixaban who presented with six days history of dyspnoea and peripheral oedema [8]. Cases of right heart thrombus have been reported in patients with nephrotic syndrome, patent foramen ovale, and COVID 19 infection however our patient did not have evidence of any of these [9,10,11,12]. Our patient had both doses of AstraZeneca vaccine that has been linked to clots in patients however there is no reported case of RV thrombus associated with it.

A study based on 12441 patients with pulmonary embolism from the Registro Informatizado de la Enfermedad TromboEmbólica (RIETE) registry and baseline echocardiogram reported the prevalence of

RHT as 2.6% [13]. The risk factors identified to increase the RHT were younger age, previous bleeding, congestive heart failure, cancer, syncope, systolic blood pressure <100 mmHg, and arterial oxyhaemoglobin saturation <90%. This study also reported that patients with RHT were significantly more likely to die from any cause and pulmonary embolism during follow-up. The outcome was worse for patients with RHT compared to those without it.

The Right Heart Thrombi European Registry, a prospective, international study reported the prognostic significance of RHT characteristics in patients with acute pulmonary embolism and Its main finding was that the clinical prognostic scores such as “simplified pulmonary embolism severity index” and “shock index” and the hemodynamic status at the time of pulmonary embolism diagnosis predicted mortality, while the RHT characteristics did not. In this study, a subgroup of patients with intermediate-risk pulmonary embolism and right ventricular strain had increased mortality risk compared with those without RHT [14].

The Moderate Pulmonary Embolism Treated with Thrombolysis trial (MOPETT trial) was another randomized control trial of thrombolytic versus anticoagulation in moderate PE. This demonstrated increased survival and reduction in PA pressure in patients receiving half-dose thrombolytic with no increased bleeding risk. This patient was discussed with a Respiratory consultant for consideration of thrombolysis however they advised against thrombolysis including half-dose thrombolysis [15].

Learning Points/Take Home Messages

- There are currently no evidence-based guidelines for the treatment of thrombus in transit associated with extensive PE and right ventricular thrombus.
- Type A thrombi are morphologically serpiginous, mobile and associated with severe PE. Thrombolysis may be indicated in this subgroup based on the patient’s hemodynamic status and in the absence of contraindications.
- Echocardiogram can be useful in the early detection of RV thrombus to assess for PFO and facilitate the treatment decision.
- Apixaban has been proven to be non-inferior to warfarin and LMWH in patients with RVT and massive pulmonary embolism.
- Patients with younger age, heart failure, cancer, recent bleeding, syncope, hypotension and hypoxemia were associated with an increased risk of RHT and carry a worse prognosis.

Patient's Perspective

It was a scary moment for me and initially I thought that I will not survive. I was very anxious about what was happening to me. I am very pleased with the care that I received at the hospital and the staff was excellent. I was kept informed all the time about my investigations and management plan and I felt much better in the end. These were probably the scariest 3 weeks I had as I could not get my head around it. I am very fit and healthy, and this was all very unexpected for me. I would rate the care I received at the hospital as excellent, and everyone was very polite and reassuring.

References

1. Nalawade D, Nawale J, Chaurasia A, Borikar N. "A case of mobile right ventricular thrombus with massive pulmonary thromboembolism". *Journal of the Practice of Cardiovascular Sciences*. 2017;3(3):178.
2. Devrey E, Bax J, Poldermans D, Vanderwall E, Holman E. "Mobile right heart thrombus and massive pulmonary embolism". *European Journal of Echocardiography*. 2007;8(3):229-231.
3. Rose P, Punjabi N, Pearse D. "Treatment of Right Heart Thromboemboli". *Chest*. 2002;121(3):806-814.
4. Nkoke C, Faucher O, Camus L, Florck L. "Free Floating Right Heart Thrombus Associated with Acute Pulmonary Embolism: An Unsettled Therapeutic Difficulty". *Case Reports in Cardiology*. Page 8 of 10 2015;2015:1-4.
5. Kronik G. "The European cooperative study on the clinical significance of right heart thrombi". *Eur Heart J* 1989;10:1046-59
6. Chartier L, Béra J, Delomez M, Asseman P, Beregi JP, Bauchart JJ, et al. "Free-floating thrombi in the right heart: Diagnosis, management, and prognostic indexes in 38 consecutive patients". *Circulation* 1999;99:2779-83
7. Pierre-Justin G, Pierard LA. "Management of mobile right heart thrombi: A prospective series". *Int J Cardiol* 2005;99:381-8.
8. MAHMOUD Z, AYAH O, JOSEPH J, AIN D. "Right Atrial Thrombus In Transit With Presumed Massive Pulmonary Embolism Treated With Apixaban". *Chest*. 2018;154(4):74A-75A.
9. Lempp S, Schwenger V. "Isolated right ventricular thrombus in an adult patient with nephrotic syndrome: a case report". *Journal of Medical Case Reports*. 2017;11(1).

10. Saunders H, Al Khalifa A, Espinosa A, Jain M. “Massive pulmonary embolism and thrombus-intransit via a patent foramen ovale: a case report of successful use of extracorporeal membrane oxygenation to manage post-embolectomy severe right ventricular dysfunction”. *European Heart Journal - Case Reports*. 2021;5(4).
11. Shepherd F, White-Stern A, Rahaman O, Kurian D, Simon K. “Saddle Pulmonary Embolism with Thrombus in Transit across a Patent Foramen Ovale”. *Case Reports in Cardiology*. 2017;2017:1-4.
12. Theetha Kariyanna P, A. Hossain N, Jayarangaiah A, A. Hossain N, Uppin V, Hegde S et al. “Thrombus in Transit and Impending Pulmonary Embolism Detected on POCUS in a Patient with COVID-19 Pneumonia”. *American Journal of Medical Case Reports*. 2020;8(8):225-228.
13. Barrios D, Rosa-Salazar V, Jiménez D, Morillo R, Muriel A, del Toro J et al. “Right heart thrombi in pulmonary embolism”. *European Respiratory Journal*. 2016;48(5):1377-1385.
14. Koc M, Kostrubiec M, Elikowski W, et al. “Outcome of patients with right heart thrombi: Right Heart Thrombi European Registry”. *Eur Respir J* 2016; 47: 869–87.
15. Sharifi M, Bay C, Skrocki L, Rahimi F, Mehdipour M. “Moderate Pulmonary Embolism Treated With Thrombolysis (from the “MOPETT” Trial)”. *The American Journal of Cardiology*. 2013;111(2):273-277.